arXiv:cond-mat/0109188v1 [cond-mat.stat-mech] 11 Sep 2001

Dynamic Transitions in Pure Ising Magnets under Pulsed and Oscillating Fields

Bikas K. Chakrabarti[1,2] and Arkajyoti Misra[2]

1st February 2008

[1]Saha Institute of Nuclear Physics, 1/AF Bidhan Nagar, Kolkata 700064, India.

[2]Theoretische Physik, Gerhard-Mercator-Universität Duisburg, D-47048, Germany.

Abstract

Response of pure Ising systems to time-dependent external magnetic fields, like pulsed and oscillating fields, are discussed and compared here. Because of the two time scales involved, namely the thermodynamic relaxation time of the system and the pulse width or the time period of the external field, dynamically broken symmetric phases appear spontaneously when both become comparable. A particularly simple case is that of an Ising ferromagnet below its static critical temperature, when it is perturbed for a short duration by a pulsed magnetic field competing with the existing order in the system. If the field strength and duration is more than the threshold (dependent on the temperature), the system, and consequently the magnetization, switches from one minimum to the other of the static free energy. This magnetization reversal transition here shows intriguing dynamic transition behaviour, similar to those for oscillating fields. Monte Carlo studies for such dynamic transitions are discussed and compared with the mean field results for the same and the Monte Carlo results for the oscillating field case. In particular, we discuss about the Monte Carlo results for the fluctuations and their growth behaviour near this magnetization reversal (dynamic) transition point.

1 Introduction

The response of pure ferromagnetic Ising systems under the action of time-dependent external magnetic fields are being studied intensively these days. A whole new class of dynamic phase transitions emerged from the study of such driven spin systems under different time dependences of the driving field. Originally it was seen that when a pure Ising system is driven by an oscillating external field, it settles to a dynamic steady state with spontaneously broken symmetry when the time period of the external field becomes smaller compared to the system's relaxation time. In this steady state, the period averaged magnetization acquires a net non-zero value, while the period averaged magnetic field remains zero [1, 2, 3, 4]. The universality class of this dynamic transition has recently been established, from extensive Monte Carlo studies, to be the same as that of the equilibrium transition in the pure Ising system [4, 5]. Later, efforts were made to investigate and analyse the response of such systems under a pulsed magnetic field which, for a short duration, either favours the existing magnetic order of the system (positive pulse) or competes with it (negative pulse). The results of the positive pulse case were analyzed by extending appropriately the finite size scaling technique to this finite time-window case, and it did not indicate any new phase transition or thermodynamic scale [6]. However, the negative pulse case indicated a new dynamic transition and thermodynamic scales, as the appropriate pulse width and strength just manages to help the system overcome the "double-well free-energy barrier" and the magnetization reversal transition occurs [7, 8, 9].

In the next section we briefly introduce the model and the magnetization reversal transition in this negative pulse case. We also briefly summarize here a few analytic results obtained earlier in the mean field limit. In section 3, we report Monte Carlo results for the fluctuations near this magnetization reversal transition in two dimensional Ising systems. In the concluding section, we analyse these observations and also compare them with those obtained for the Ising model under oscillating field.

2 Model

The model we consider here is the Ising model with nearest-neighbour interaction under a time-dependent external magnetic field, described by the Hamiltonian

$$H = -\frac{J}{2}\sum_{[ij]} S_i S_j - h(t)\sum_i S_i. \tag{1}$$

Here, J denotes the strength of the cooperative interaction between nearest neighbour spins put under a time-dependent field $h(t)$. S_i and S_j denote the spins at lattice sites i and j respectively and the nearest-neighbour spin pairs are denoted by [...]. We consider such a system at temperatures T below its static critical temperature T_c. The external field is applied at $t = t_0$ after the system is brought to equilibrium, having an equilibrium magnetization m_0 (T). The external field $h(t)$ has a time dependence given by $h(t) = -h_p$ for $t_0 \leq t \leq t_0 + \Delta t$ and $h(t) = 0$ outside this range. Typical responses of the time-dependent magnetization $m(t)$ of the system under different $h(t)$ are indicated in Fig. 1. For appropriate combinations of h_p and Δt, at a fixed T, a magnetization reversal transition occurs when the magnetization of the system eventually switches to $-m_0$. It may noted that a magnetization reversal phenomenon trivially occurs in the limit $\Delta t \to \infty$ for any non-vanishing value of h_p at any temperature $T < T_c$. This is however a limiting case of the transition, which is studied here for any finite value of Δt . Also, in the case of magnetization reversal transition, the magnetization reversal need not occur during the presence of the external pulsed field. In fact, the time required for the system to reach its final equilibrium state diverges as the transition point is approached [7].

A qualitative understanding for these diverging time and also length scales at this magnetization reversal transition point can be easily understood from a mean field study of this problem [7, 8]. Using mean field approximation for the dynamics, we can write the equation of motion for the average magnetization $m_i (=< S_i >)$ as

$$\frac{\mathrm{d}m_i}{\mathrm{d}t} = -m_i + \tanh\left(\frac{J\sum_j m_j + h(t)}{T}\right), \tag{2}$$

where $< ... >$ denotes the thermodynamic (ensemble) average. This equation may be linearized near the magnetization reversal transition point and for uniform magnetization one gets

$$m(t) = m_0 - \left(\frac{h_p}{\Delta T} - m_0\right)\left[\exp\left\{\frac{\Delta T}{T}(t - t_0)\right\}\right] \tag{3}$$

as a solution of Eq. (2). Here $\Delta T = T_c^{\mathrm{mf}} - T$, where $T_c^{\mathrm{mf}} \equiv J(q = 0)$ is the critical temperature in the mean field approximation and $J(q)$ is the Fourier transform of the interaction J. From the solution (3) the value of m_0 is always restricted to be less than $h_p/\Delta T$ in order to make the linear approximation valid. Hence during the presence of the field, $m(t)$ will decrease in its magnitude from $m(t_0) \equiv m_0$ to $m(t_0 + \Delta t) \equiv m_w$ at the time of withdrawal of the pulse. Since there is no fluctuation here, the magnetization will relax back to its original value m_0 if m_w is positive or relax to a value $-m_0$ if m_w is negative. This can be easily seen by solving eq. (2) for $t > t_0 + \Delta t$ where $h(t) = 0$, using m_w as the initial value of the magnetization. In this time region one gets a Debye type relaxation of magnetization to its final equilibrium value, with a relaxation time [7]

$$\tau \sim \left(\frac{1}{T_c^{\mathrm{mf}} - T}\right) \ln\left(\left|\frac{m_0}{m_w}\right|\right).$$

It diverges at the magnetization reversal transition point, where m_w vanishes. The prefactor, giving the divergence of τ at the static mean field transition temperature, is responsible for the usual critical slowing down phenomenon at the static transition point ($h = 0$). The other factor gives the diverging time scale of interest here, at any temperature below the static transition temperature, where magnetization reversal occurs or m_w vanishes due to the appropriate combination of the external field pulse strength and duration. A similar solution for the Fourier transform of the linearized version of the equation (2), with an additional derivative with respect to the field h for the susceptibility $\chi(q)$, gives [8, 9]

$$\chi(q) \sim \exp\left(-q^2\xi^2\right),$$

where

$$\xi \sim \left[\frac{1}{(T_c^{\mathrm{mf}} - T)} \ln\left(\frac{1}{|m_w|}\right)\right]^{\frac{1}{2}}.$$

Here also, the prefactor in ξ gives the usual divergence at T_c^{mf} , while the other factor gives the divergence of interest here at the magnetization reversal transition point. Such divergences of τ and ξ across the dynamic transition phase boundary which gives the optimal combination of h_p and Δt at any particular T, are also seen [9] in the Monte Carlo simulations.

We believe, a similar transition and the associated length and time scales are responsible in an essential way for the spontaneous dynamic symmetry breaking transition in the Ising systems under oscillating fields [1, 5]. The growing time scale induced by the periodically changing field competing with the existing order in the system at that point, locks the system dynamically in a broken symmetric phase. In order to compare these scenario, we have studied here the growing fluctuations in the order parameter m_w and the internal energy E, as one approaches the magnetization reversal phase boundary in a Monte Carlo study. We find that the fluctuations in both these quantities diverge very prominently at the magnetization reversal phase boundary. We also indicate the behaviour of the derivatives $(\mathrm{d}m_w/\mathrm{d}h_p)$ and $(\mathrm{d}E/\mathrm{d}T)$, which show some peak across the same phase boundary. We compare our results with those reported for the dynamic transition under oscillating field [10, 5].

3 Monte Carlo study

We study here the behaviour of the fluctuations near the magnetization reversal transition in a ferromagnetic Ising system on square lattice of size L, under negative or competing field pulses, using Monte Carlo simulations with single spin flip Glauber dynamics. Working at temperatures T below the static critical temperature T_c ($\simeq 2.27$ in units of J), the system is brought to the equilibrium state by evolving the system without any external field, from an initial configuration (say all spins up). The equilibrium relaxation time required for this system to evolve to its equilibrium state with magnetization m_0 is noted. The field pulse of strength $-h_p$ for a duration Δt (measured in Monte Carlo steps or MCS), competing with this order, is applied at $t = t_0$ only after the system has reached its equilibrium. The magnetization $m(t)$ starts decreasing (from its equilibrium value m_0) due to the effect of the competing field $-h_p$ during the time period Δt. The average value of the magnetization m_w at the time of the withdrawal of the pulse is noted. The phase boundary of this dynamic transition gives the appropriate combination of h_p and Δt that induces the magnetization reversal by bringing $m(t)$ from $m(t_0) = m_0$ to $m(t_0 + \Delta t) \equiv m_w = 0$.

Therefore across the phase boundary m_w changes its sign [7]. Obviously the phase boundary changes with T. We study the behaviour of various thermodynamic quantities across these phase boundaries. The thermodynamic quantities are averaged over different initial configurations of the system. We note the fluctuations over these average quantities as well. In our simulations we have considered three different system sizes, viz., L =256, 512 and 1024 for all our studies reported here. The typical number of averages is 200 for $L = 1024$ and 1000 for $L = 256$.

In Fig. 2 we show the growth of fluctuations $\Delta m_w = L^2(< m_w^2 > - < m_w >^2)$ in m_w across the phase boundary at two different points, both driven by the pulse strength h_p. The insets here show the corresponding variations of the pulse-withdrawal-time magnetization m_w, which crosses zero at the phase boundary. Figure 3 shows the growth of fluctuations $\Delta E_{\text{tot}} = L^2(< E_{\text{tot}}^2 > - < E_{\text{tot}} >^2)$ in the average total energy E_{tot} per spin pair (in units of J) across the phase boundary, again at two points, both driven by T. The insets here again show the corresponding m_w variations. Both these fluctuations Δm_w and ΔE_{tot} clearly diverge at the phase boundary, in the large system size limit. Fig. 4 shows the variation of the fluctuation $\Delta E_{\text{coop}} = L^2(< E_{\text{coop}}^2 > - < E_{\text{coop}} >^2)$ in the cooperative part E_{coop} (excluding the field contribution) of the total energy E_{tot} for the same parameter values as those of Fig. 3(a). Although the fluctuation in E_{coop} also grows in general around the transition point, a sharp dip is observed there. This dip is identified to be due to the sharp growth of E_{coop} at the transition point, as shown in the inset.

In Fig. 5, we show the behaviour of the derivatives $\mathrm{d}m_w/\mathrm{d}h_p$ (in (a)) and $\mathrm{d}E_{\text{tot}}/\mathrm{d}T$ (in (b)) across the phase boundary at two different points. Unlike in the case of the dynamic transition under oscillating fields [10], we do not observe any divergence in these quantities at the transition point. This is because the curves are independent of system size and hence the peak values do not grow with L. However, the peaks become sharper at lower temperatures (higher values of h_p) as the system approaches the tricritical point occurring on the phase boundary [9].

4 Concluding remarks

We have studied here the growth of fluctuations and their system size dependence across the dynamic magnetization reversal phase boundary. This transition can be driven by the field pulse strength h_p or its duration Δt or even by varying T in the presence of a fixed field pulse but always keeping it below T_c. 'Critical slowing down' and the divergence of length scales etc at this dynamic phase boundary have already been demonstrated earlier [9], using both mean field approximation and the Monte Carlo simulations in two and three dimensions. Here we have shown the Monte Carlo results for the growth of fluctuations, both in the field-withdrawal-time magnetization m_w and in the total energy E_{tot} , across the dynamic phase transition boundary in two dimension. The growth of these fluctuations with the system size L clearly indicates a divergence at the phase boundary. It may also be noted that although we have shown the fluctuations for one specific driving parameter, e.g. Δm_w with h_p and ΔE_{tot} with T, similar fluctuations in both the quantities are seen for other driving parameters as well. All these clearly indicate a genuine dynamic phase transition at the magnetization reversal point of an Ising system under a competing field pulse.

Acknowledgement: BKC is grateful to K. D. Usadel for his kind hospitality in the Theoretical Physics Group, University of Duisburg, where a part of the work was done and the manuscript was finalized. A part of the work has been supported by the Deutsche Forschungsgemeinschaft through SFB 491 and Graduiertenkolleg 277.

References

[1] B. K. Chakrabarti and A. Acharyya, Rev. Mod. Phys. **71** (1999) 847

[2] W. S. Lo and R. A. Pelcovits, Phys. Rev. A **42** (1990) 7471

[3] M. Acharyya and B. K. Chakrabarti, Phys. Rev. B **52** (1995) 6550

[4] S. W. Sides, P. A. Rikvold and M. A. Novotny, Phys. Rev. Lett. **81** (1998) 834; Phys. Rev. E **59** (1999) 2710

[5] G. Korniss, C. J. White, P. A. Rikvold and M. A. Novotny, Phys. Rev. E **63** (2001) 016120

[6] M. Acharyya, J. K. Bhattacharjee and B. K. Chakrabarti, Phys. Rev. E **55** (1997) 2392

[7] A. Misra and B. K. Chakrabarti, Phys. Rev. E **58** (1998) 4277

[8] R. B. Stinchcombe, A. Misra and B. K. Chakrabarti, Phys. Rev. E **59** (1999) R4717

[9] A. Misra and B. K. Chakrabarti, J. Phys. A **33** (2000) 4249; Europhys. Lett. **52** (2000) 311

[10] M. Acharyya, Phys. Rev. E **56** (1997) 2407

Figure Captions:

Fig. 1 Typical time variation of the response magnetizations $m(t)$ for two different field pulses $h(t)$ with different strengths h_p, but same width Δt and at same temperature T, are shown. The quantities of interest to characterize the response magnetization are also indicated.

Fig. 2 Growth of fluctuations Δm_w near the transition, driven by h_p, are shown for two different points on the phase boundary. (a) for $T = 1.0$ and $\Delta t = 12$ MCS, and (b) for $T = 1.50$ and $\Delta t = 2$ MCS. The insets show the corresponding m_w variations; $m_w = 0$ at the transition point.

Fig. 3 Growth of fluctuations $\Delta E_{\rm tot}$ near the transition, driven by T , are shown for two different points on the phase boundary. (a) for $h_p = 1.75$ and $\Delta t = 10$ MCS, and (b) for $h_p = 1.25$ and $\Delta t = 10$ MCS. The insets show the corresponding m_w variations; $m_w = 0$ at the transition point.

Fig. 4 Growth of fluctuations $\Delta E_{\rm coop}$ near the transition point at $h_p = 1.75$ and Δt =10 MCS. The insets show the (rather sharp) variations of $< E_{\rm coop} >$ and $< E^2_{\rm coop} >$ at the transition point, giving rise to the dip there in $\Delta E_{\rm coop}$.

Fig. 5 (a) The variation of $\mathrm{d}m_w/\mathrm{d}h_p$ across two different points on the phase boundary: for T =1.50 and $\Delta t = 2$ MCS (open symbol) and $T = 1.00$ and Δt $= 12$ MCS (solid symbol); (b) The variation of $\mathrm{d}E_{\rm tot}/\mathrm{d}T$ across two different points on the phase boundary: for $h_p = 1.25$ and Δt =10 MCS (open symbol) and $h_p = 1.75$ and $\Delta t = 10$ (solid symbol).



Fig. 1 Chakrabarti et al

Fig. 2 Chakrabarti et al



Fig. 2 Chakrabarti et al



Fig. 3 Chakrabarti et al



Fig. 3 Chakrabarti et al



Fig. 4 Chakrabarti et al





Fig. 5 Chakrabarti et al



Fig. 5 Chakrabarti et al